

Duality and Integrability of Two Dimensional String Effective Action

Ashok Das^a, J. Maharana^{b*} and A. Melikyan^a

^a *Department of Physics and Astronomy, University of Rochester, NY 14627-0171, USA*

^b *Theory Division, KEK, Tsukuba, Ibaraki 3050801, Japan*

We present a prescription for constructing the monodromy matrix, $\widehat{\mathcal{M}}(\omega)$, for $O(d, d)$ invariant string effective actions and derive its transformation properties under the T -duality group. This allows us to construct $\widehat{\mathcal{M}}(\omega)$ for new backgrounds, starting from known ones, which are related by T -duality. As an application, we derive the monodromy matrix for the exactly solvable Nappi-Witten model, both when $B = 0$ and $B \neq 0$.

The symmetry structure of string theories is one of its most fascinating features. In particular, duality symmetries have played a very important role in our understanding of string dynamics [1–3]. Whereas T -duality properties can be subjected to tests within the perturbative frame work, the S -duality symmetry explores nonperturbative attributes of string theories. The tree level string effective action, compactified on a d -dimensional torus, T^d , is known to be invariant under the noncompact global symmetry group $O(d, d)$. Indeed, this symmetry property has been exploited, in the past, to generate new string vacua from known solutions.

The dimensionally reduced string effective action, in two space-time dimensions, is known to possess enlarged symmetries, which has been explored by several authors [4–7]. The effective action is described by a nonlinear σ -model, defined over a coset, which is coupled to gravity. In the past, two dimensional models, derived from dimensional reduction of higher dimensional Einstein gravity as well as supergravity theories, have been studied to bring out their integrability properties [8–10]. In this context, one may recall that the construction of the monodromy matrix turns out to be one of the principal objectives in the study of integrable systems [11,12]. Furthermore, the integrability properties of such models have been studied in the past in some detail from different perspectives. Recently, we have investigated the scattering of plane fronted waves, in four dimensions, which correspond to massless states of closed strings [13]. The scattering process is described by an effective two dimensional action due to the presence of isometries along the directions transverse to the axis of collision. Therefore, the reduced action is endowed with an $O(2, 2)$ symmetry, as is obvious from the preceding discussion. We were able to derive the monodromy matrix for such a scattering process for the specific background configurations involved and had established an intimate connection between the integrability properties of the model and the underlying $O(2, 2)$ symmetry.

The purpose of this note is to derive the monodromy matrix, $\widehat{\mathcal{M}}(\omega)$, for a 2-dimensional string effective action, obtained from a D -dimensional effective action, which is compactified on T^d and thus enjoys isometries along $d = D - 2$ spatial directions. We provide the prescription for the construction of the monodromy matrix, in this case, and derive the transformation property of $\widehat{\mathcal{M}}(\omega)$ under $O(d, d)$ transformations. In this process, we are able to synthesize the classical integrability properties and the duality symmetry of string theory in a novel way. Furthermore, as an illustrative example, we construct the monodromy matrix for the well known Nappi-Witten model [14], both for $B = 0$ and $B \neq 0$, and demonstrate explicitly the transformation of $\widehat{\mathcal{M}}(\omega)$ under T -duality. Namely, as is well known, one can start from background configurations where the NS-NS field $B_{ij} = 0$ and generate new backgrounds with $B_{ij} \neq 0$ through a T -duality rotation. Correspondingly, we can construct $\widehat{\mathcal{M}}^{(B=0)}(\omega)$ explicitly and generate $\widehat{\mathcal{M}}^{(B)}(\omega)$ through a T -duality rotation. Alternatively, we can also construct the $\widehat{\mathcal{M}}^{(B)}(\omega)$ directly and demonstrate that the two methods indeed lead to identical results.

Our starting point is the D -dimensional tree level string effective action

$$\hat{S} = \int d^D x \sqrt{-\hat{G}} \left(R_{\hat{G}} + (\partial\hat{\phi})^2 - \frac{1}{12} \hat{H}^2 \right) \quad (1)$$

*Permanent address: Institute of Physics, Bhubaneswar 751005, India

where $\hat{G}_{\hat{\mu}\hat{\nu}}$ is the metric in D -dimensions in the string frame, $\hat{G}$ its determinant, $R_{\hat{G}}$ the corresponding scalar curvature, $\hat{\phi}$ is the dilaton and $\hat{H} = d\hat{B}$. When compactified on T^d , the reduced action takes the following form [15]:

$$S = \int dx^0 dx^1 \sqrt{-g} e^{-\bar{\phi}} \left(R + (\partial\bar{\phi})^2 + \frac{1}{8} \text{Tr}(\partial_\alpha M^{-1} \partial^\alpha M) \right) \quad (2)$$

where $g_{\alpha\beta}$ with $\alpha, \beta = 0, 1$ is the two dimensional space-time metric, and R is the corresponding scalar curvature. The shifted dilaton is $\bar{\phi} = \phi - \frac{1}{2} \log \det G$, with G_{ij} the metric corresponding to the transverse coordinates $x^i, i = 2, 3, \dots, D-1$. M is a $2d \times 2d$ symmetric matrix belonging to $O(d, d)$,

$$M = \begin{pmatrix} G^{-1} & -G^{-1}B \\ BG^{-1} & G - BG^{-1}B \end{pmatrix} \quad (3)$$

where B represents the moduli coming from the reduction of the B -field in D space-time dimensions. The symmetric nature of M is evident since G is symmetric and B is anti-symmetric. In general, there will be additional terms in (2) corresponding to d Abelian gauge fields coming from the original metric and another set of d gauge fields coming from the anti-symmetric tensor $B_{\hat{\mu}\hat{\nu}}$ as a result of dimensional reduction [15] ($d = D - 2$). Furthermore, there would also have been the field strength of the two dimensional tensor field $B_{\alpha\beta}$. Since we are effectively in two space-time dimensions, we have dropped the gauge field terms and, in the same spirit, have not kept the field strength of $B_{\alpha\beta}$, which can be removed if it depends only on the coordinates x^0 and x^1 .

The action (2) is invariant under the global $O(d, d)$ transformations,

$$g_{\alpha\beta} \rightarrow g_{\alpha\beta}, \quad \bar{\phi} \rightarrow \bar{\phi}, \quad M \rightarrow \Omega^T M \Omega \quad (4)$$

where $\Omega \in O(d, d)$ is the transformation matrix, which leaves the $O(d, d)$ metric, $\eta = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$, invariant ($\mathbf{1}$ is the d -dimensional unit matrix). It is convenient to introduce the lower triangular matrix

$$V = \begin{pmatrix} E^{-1} & 0 \\ BE^{-1} & E^T \end{pmatrix} \quad (5)$$

where E is the vielbein with $(EE^T)_{ij} = G_{ij}$. It is easy to check that $V \in O(d, d)$ and that $M = VV^T$ so that under $O(d, d)$ transformations, $V \rightarrow \Omega^T V$. In fact, V belongs to the coset $\frac{O(d, d)}{O(d) \times O(d)}$.

We may decompose the current as

$$V^{-1} \partial_\alpha V = P_\alpha + Q_\alpha \quad (6)$$

where $Q_\alpha \in O(d) \times O(d)$, the Lie algebra for the maximally compact subgroup and P_α belongs to the complement. It follows from the property of the coset under study that $P_\alpha^T = P_\alpha$ and $Q_\alpha^T = -Q_\alpha$ and it is straightforward to check that the last term of (2) can be expressed as

$$\text{Tr}(\partial_\alpha M \partial^\alpha M^{-1}) = -4 \text{Tr} g^{\alpha\beta} P_\alpha P_\beta \quad (7)$$

Let us note that, under a global $O(d, d)$ rotation $V \rightarrow \Omega^T V$, the current in (6) is invariant, while under a local $O(d) \times O(d)$ transformation $V \rightarrow Vh(x)$, P_α and Q_α transform as

$$P_\alpha \rightarrow h^{-1}(x) P_\alpha h(x), \quad Q_\alpha \rightarrow h^{-1}(x) Q_\alpha h(x) + h^{-1}(x) \partial_\alpha h(x) \quad (8)$$

which leaves (7) invariant.

A few comments are in order at this stage. We are dealing with a 2-dimensional σ -model [4,6,7,16] coupled to gravity and, in addition, the action (2) contains a shifted dilaton field, $\bar{\phi}$. We may rescale the metric $g_{\alpha\beta} \rightarrow e^{\bar{\phi}} g_{\alpha\beta}$, which will eliminate the kinetic energy term of $\bar{\phi}$. Recall that in a curved space, the spectral parameter becomes space-time dependent in the zero curvature formulation of the sigma model, unlike in flat space, where the spectral parameter is a constant. Let $\hat{V}(x, t)$, where t is the space-time dependent spectral parameter, be the one parameter family of matrices such that $\hat{V}(x, t=0) = V(x)$ and

$$\hat{V}^{-1}(x, t) \partial_\alpha \hat{V}(x, t) = Q_\alpha + \frac{1+t^2}{1-t^2} P_\alpha + \frac{2t}{1-t^2} \epsilon_{\alpha\beta} P^\beta \quad (9)$$

Then, it is easy to check that the integrability condition

$$\partial_\alpha(\widehat{V}^{-1}\partial_\beta\widehat{V}) - \partial_\beta(\widehat{V}^{-1}\partial_\alpha\widehat{V}) + [\widehat{V}^{-1}\partial_\alpha\widehat{V}, \widehat{V}^{-1}\partial_\beta\widehat{V}] = 0 \quad (10)$$

leads to the dynamical equations of the theory, provided the spectral parameter is related to the shifted dilaton through the equation

$$\partial_\alpha t = -\frac{1}{2}\epsilon_{\alpha\beta}\partial^\beta \left(e^{-\bar{\phi}} \left(t + \frac{1}{t} \right) \right) \quad (11)$$

The solution to this equation has the form

$$t(x) = \frac{\sqrt{\omega + \rho_+(x^+)} - \sqrt{\omega - \rho_-(x^-)}}{\sqrt{\omega + \rho_+(x^+)} + \sqrt{\omega - \rho_-(x^-)}} \quad (12)$$

where we have defined $\rho = e^{-\bar{\phi}} = \rho_+(x^+) + \rho_-(x^-)$ for later use and ω is the constant of integration (which can be thought of as a global, constant spectral parameter). The light cone coordinates are defined to be $x^+ = \frac{1}{\sqrt{2}}(x^0 + x^1)$ and $x^- = \frac{1}{\sqrt{2}}(x^0 - x^1)$.

The next important step is to solve for the monodromy matrix, $\widehat{\mathcal{M}}(\omega)$. The linear system (9) is known to be invariant under a generalization of the symmetric space automorphism, τ^∞ , defined through

$$\tau^\infty \widehat{V}(x, t) = \left(\widehat{V}^{-1} \right)^T \left(x, \frac{1}{t} \right) \quad (13)$$

This relation has the following action on the elements of the Lie algebra, $Q_\alpha \rightarrow Q_\alpha$, $P_\alpha \rightarrow -P_\alpha$ and $t \rightarrow \frac{1}{t}$. The monodromy matrix, defined as,

$$\widehat{\mathcal{M}}(\omega) = \widehat{V}(x, t) \tau^\infty \widehat{V}^{-1}(x, t) = \widehat{V}(x, t) \widehat{V}^T(x, \frac{1}{t}) \quad (14)$$

is space-time independent and plays a cardinal role in the study of integrable models. The reconstruction of solutions and properties of integrability are all deeply encoded in $\widehat{\mathcal{M}}(\omega)$. For example, if we were to use a solution of the $\bar{\phi}$ equation as input, then we can determine t from its relation with the dilaton and eventually obtain $\widehat{V}(x, t)$.

Now we want to present a crucial observation which enables us to derive the transformation property of $\widehat{\mathcal{M}}(\omega)$ under T -duality. Note that the one parameter family of matrices $\widehat{V}(x, t) \in \frac{O(d, d)}{O(d) \times O(d)}$ much like $V(x) = \widehat{V}(x, t = 0)$. Therefore, under a global $O(d, d)$ rotation, $\widehat{V}(x, t) \rightarrow \Omega^T \widehat{V}(x, t)$ and the monodromy matrix transforms as

$$\widehat{\mathcal{M}}(\omega) \rightarrow \Omega^T \widehat{\mathcal{M}}(\omega) \Omega \quad (15)$$

Furthermore, although under a local $O(d) \times O(d)$ transformation $\widehat{V}(x, t) \rightarrow \widehat{V}(x, t) h(x)$, the only local transformations, which will preserve the global nature of the monodromy matrix are the ones which do not depend on the spectral parameter explicitly. This is very much like what we already know for the case $t = 0$, namely, although $V \rightarrow \Omega^T V h(x)$ under combined $O(d, d)$ and $O(d) \times O(d)$ transformations, $M = V V^T$ undergoes only a global $O(d, d)$ rotation. In other words, the monodromy matrix, $\widehat{\mathcal{M}}(\omega)$, is only affected by the global $O(d, d)$ rotation, as is the case with the M -matrix. Thus, we are able to synthesize the integrability properties of the two dimensional string effective action with its T -duality properties. Indeed, this relation can be explicitly checked in specific examples (as we will show shortly) and can be used as a powerful tool in determining solutions. Let us derive the transformation properties of $\widehat{\mathcal{M}}$ under infinitesimal $O(d, d)$ transformations. We may express $\widehat{\mathcal{M}}$ as

$$\widehat{\mathcal{M}} = \begin{pmatrix} \widehat{\mathcal{M}}_{11} & \widehat{\mathcal{M}}_{12} \\ \widehat{\mathcal{M}}_{21} & \widehat{\mathcal{M}}_{22} \end{pmatrix} \quad (16)$$

where each element is a $d \times d$ matrix. The matrix Ω assumes the following form

$$\Omega = \begin{pmatrix} 1 + X & Y \\ Z & 1 + W \end{pmatrix} \quad (17)$$

where the infinitesimal parameters of the transformation are required to satisfy, $Y^T = -Y$, $Z^T = -Z$ and $W = -X^T$. Under (17), the elements of (16) transform as

$$\begin{aligned}
\delta \widehat{\mathcal{M}}_{11} &= \widehat{\mathcal{M}}_{11} X + X^T \widehat{\mathcal{M}}_{11} - Z \widehat{\mathcal{M}}_{12} + \widehat{\mathcal{M}}_{12} Z \\
\delta \widehat{\mathcal{M}}_{12} &= \widehat{\mathcal{M}}_{11} Y + X^T \widehat{\mathcal{M}}_{12} - Z \widehat{\mathcal{M}}_{22} - \widehat{\mathcal{M}}_{12} X^T \\
\delta \widehat{\mathcal{M}}_{21} &= -Y \widehat{\mathcal{M}}_{11} - X \widehat{\mathcal{M}}_{21} + \mathcal{M}_{21} X + \widehat{\mathcal{M}}_{22} Z \\
\delta \widehat{\mathcal{M}}_{22} &= \widehat{\mathcal{M}}_{21} Y - Y \mathcal{M}_{12} - X \widehat{\mathcal{M}}_{22} - \widehat{\mathcal{M}}_{22} X^T
\end{aligned} \tag{18}$$

Since, we can obtain $\widehat{\mathcal{M}}(\omega)$ for the case when $B \neq 0$ starting from a vanishing B configuration, let us derive $\widehat{\mathcal{M}}(\omega)$ for the case when $B = 0$, E is diagonal so that G_{ij} is also diagonal. It is easy to check that $Q_\alpha = 0$ for this choice of backgrounds and the relevant equation for determining $\widehat{V}^{(B=0)}$ are given by

$$(\widehat{V}^{(B=0)})^{-1} \partial_+ \widehat{V}^{(B=0)} = \frac{1-t}{1+t} P_+ \quad \text{and} \quad (\widehat{V}^{(B=0)})^{-1} \partial_- \widehat{V}^{(B=0)} = \frac{1+t}{1-t} P_- \tag{19}$$

and t satisfies (12). The factorizability property of $\widehat{\mathcal{M}}(\omega)$, namely (14), together with the assumption that it has only isolated single poles determines that the monodromy matrix (as well as the $\widehat{V}^{(B=0)}(x, t)$ matrix) will have $2d$ poles in the spectral parameter. We can check this in the following manner. Let us assume that G_{ij} has the diagonal structure

$$G = \text{diag}(e^{\lambda+\psi_1}, e^{\lambda+\psi_2}, \dots, e^{\lambda+\psi_d}) \tag{20}$$

with $\sum \psi_i = 0$, so that $\lambda = \frac{1}{d} \log \det G$ as adopted in [18]. The E -matrix is also diagonal and has the form

$$E = \text{diag}(e^{\frac{1}{2}(\lambda+\psi_1)}, e^{\frac{1}{2}(\lambda+\psi_2)}, \dots, e^{\frac{1}{2}(\lambda+\psi_d)}) \tag{21}$$

Correspondingly, the $\widehat{V}^{(B=0)}$ matrix is diagonal as well and has the form

$$\widehat{V}^{(B=0)}(x, t) = \begin{pmatrix} \overline{V}^{(B=0)}(x, t) & 0 \\ 0 & (\overline{V}^{(B=0)})^{-1}(x, t) \end{pmatrix} \tag{22}$$

Let us assume the following form for the diagonal matrix $\overline{V}^{(B=0)}(x, t) = (\overline{V}_1, \overline{V}_2, \dots, \overline{V}_d)$ with

$$\overline{V}_i = \frac{t_{d+i}}{t_i} \frac{t-t_i}{t-t_{d+i}} E_i^{-1} \tag{23}$$

which explicitly brings out the $2d$ pole structures in $\widehat{V}^{(B=0)}(x, t)$. The spectral parameters satisfy

$$\partial_\pm t = \frac{1 \mp t}{1 \pm t} \partial_\pm \ln \rho, \quad \partial_\pm t_i = \frac{1 \mp t_i}{1 \pm t_i} \partial_\pm \ln \rho \tag{24}$$

Using this, it is easy to verify that $(t_{d+i}$ and t_i have opposite signatures, following from the double valued nature of the solutions in Eq. (12))

$$\overline{V}_i^{-1} \partial_\pm \overline{V}_i = \partial_\pm \ln E_i^{-1} \mp \frac{t}{1 \pm t} \partial_\pm \ln \left(-\frac{t_i}{t_{d+i}} \right) = \frac{1 \mp t}{1 \pm t} \partial_\pm \ln E_i^{-1} \tag{25}$$

provided we identify $-\frac{t_i}{t_{d+i}} = E_i^{-2}$. In such a case, we can write

$$(\widehat{V}^{(B=0)})^{-1}(x, t) \partial_\pm \widehat{V}^{(B=0)}(x, t) = \frac{1 \mp t}{1 \pm t} \begin{pmatrix} -E^{-1} \partial_\pm E & 0 \\ 0 & E^{-1} \partial_\pm E \end{pmatrix} = \frac{1 \mp t}{1 \pm t} P_\pm \tag{26}$$

Thus, we see that, for the case at hand, we have to introduce $2d$ poles, a pair for every diagonal element E_i . Furthermore, the diagonal elements of $\overline{V}^{(B=0)}(x, t)$, in this case, are determined to be

$$\overline{V}_i(x, t) = \frac{t_{d+i}}{t_i} \frac{t-t_i}{t-t_{d+i}} E_i^{-1} = \sqrt{-\frac{t_{d+i}}{t_i}} \frac{t-t_i}{t-t_{d+i}} \tag{27}$$

It is easy to check that the global (constant) spectral parameters satisfy

$$\frac{\omega - \omega_i}{\omega - \omega_{d+i}} = \frac{t_{d+i}}{t_i} \frac{t-t_i}{t-t_{d+i}} \frac{\frac{1}{t} - t_i}{\frac{1}{t} - t_{d+i}} \tag{28}$$

Using this relation as well as the definition of the monodromy matrix, we note that we can write

$$\widehat{\mathcal{M}}^{(B=0)}(\omega) = \widehat{V}^{(B=0)}(x, t) (\widehat{V}^{(B=0)})^T(x, \frac{1}{t}) = \begin{pmatrix} \mathcal{M}(\omega) & 0 \\ 0 & \mathcal{M}^{-1}(\omega) \end{pmatrix} \quad (29)$$

where $\mathcal{M}(\omega)$ is diagonal with

$$\mathcal{M}_i(\omega) = \overline{V}_i(x, t) \overline{V}_i(x, \frac{1}{t}) = -\frac{\omega - \omega_i}{\omega - \omega_{d+i}} \quad (30)$$

The double valued relation between the global and the local spectral parameters allows us to choose $\omega_{d+i} = -\omega_i$, which leads to

$$\mathcal{M}_i(\omega) = \frac{\omega_i - \omega}{\omega_i + \omega} \quad (31)$$

This determines the monodromy matrix for a pure G background and starting from this, we can determine the monodromy matrix for a background with a non-vanishing G and B generated through an $O(d, d)$ transformation.

We shall adopt the Nappi-Witten model to illustrate these points explicitly. The Nappi-Witten model is described by the following $\frac{SU(2)}{U(1)} \times \frac{SL(2, R)}{SO(1, 1)}$ gauged WZW model [14]. The worldsheet action, in terms of holomorphic and anti-holomorphic coordinates, is:

$$S = \int d^2 z (-\partial\psi\bar{\partial}\psi + \partial s\bar{\partial}s + \partial\rho\bar{\partial}\rho G_{\rho\rho} + \partial\lambda\bar{\partial}\lambda G_{\lambda\lambda} + (\partial\rho\bar{\partial}\lambda - \partial\lambda\bar{\partial}\rho)B) \quad (32)$$

where (we will identify $x^0 = \tau$ here)

$$G_{\rho\rho} = \frac{4\cos^2\tau\cos^2x}{(1 - \cos 2\tau\cos 2x)}, \quad G_{\lambda\lambda} = \frac{4\sin^2\tau\sin^2x}{(1 - \cos 2\tau\cos 2x)}, \quad B = \frac{(\cos 2\tau - \cos 2x)}{(1 - \cos 2\tau\cos 2x)} \quad (33)$$

and $\bar{\phi} = -\ln(\sin 2x \sin 2\tau)$. This model describes a 4-dimensional cosmological scenario with graviton, dilaton and anti-symmetric tensor fields and possesses two isometries.

The backgrounds, in Eq. (33), can be generated from the ones where $B = 0$ and the metric given by:

$$ds^2 = -d\tau^2 + dx^2 + \tan^{-2}\tau dy^2 + \tan^2 x dz^2 \quad (34)$$

through an $O(2, 2)$ transformation (see [17] for a more general discussion):

$$\Omega = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \quad (35)$$

We can construct $\mathcal{M}(\omega)$ corresponding to these two vacua following the procedure described earlier (see Eqs. (27, 31))

$$\widehat{V}^{(B=0)}(x, t) = \text{diag}(\sqrt{-\frac{t_3}{t_1}} \frac{t-t_1}{t-t_3}, \sqrt{-\frac{t_4}{t_2}} \frac{t-t_2}{t-t_4}, \sqrt{-\frac{t_1}{t_3}} \frac{t-t_3}{t-t_1}, \sqrt{-\frac{t_2}{t_4}} \frac{t-t_4}{t-t_2}) \quad (36)$$

Thus, the monodromy matrix for $B = 0$ follows to have the form

$$\widehat{\mathcal{M}}^{(B=0)}(\omega) = \begin{pmatrix} \frac{\omega_1 - \omega}{\omega_1 + \omega} & 0 & 0 & 0 \\ 0 & \frac{\omega_2 - \omega}{\omega_2 + \omega} & 0 & 0 \\ 0 & 0 & \frac{\omega_1 + \omega}{\omega_1 - \omega} & 0 \\ 0 & 0 & 0 & \frac{\omega_2 + \omega}{\omega_2 - \omega} \end{pmatrix} = \begin{pmatrix} \mathcal{M}(\omega) & 0 \\ 0 & \mathcal{M}^{-1}(\omega) \end{pmatrix} \quad (37)$$

We can also compute $\widehat{V}^{(B)}(x, t)$ directly from the Nappi-Witten action which takes the form:

$$\widehat{V}^{(B)}(x, t) = \frac{1}{\sqrt{2}} \begin{pmatrix} \gamma\hat{V}_1 + \delta\hat{V}_4 & 0 & 0 & \delta\hat{V}_1 - \gamma\hat{V}_4 \\ 0 & \gamma\hat{V}_2 + \delta\hat{V}_3 & -\delta\hat{V}_2 + \gamma\hat{V}_3 & 0 \\ 0 & -\gamma\hat{V}_2 + \delta\hat{V}_3 & \delta\hat{V}_2 + \gamma\hat{V}_3 & 0 \\ \gamma\hat{V}_1 - \delta\hat{V}_4 & 0 & 0 & \delta\hat{V}_1 + \gamma\hat{V}_4 \end{pmatrix} \quad (38)$$

where $\gamma = \sqrt{\frac{1-B}{2}}$, $\delta = \sqrt{\frac{1+B}{2}}$, and $\hat{V}_i$ are the diagonal elements of $\hat{V}^{(B=0)}(x, t)$ in (36). This leads to

$$\widehat{\mathcal{M}}^{(B)}(\omega) = \frac{1}{2} \begin{pmatrix} \mathcal{M}_1 + \mathcal{M}_2^{-1} & 0 & 0 & \mathcal{M}_1 - \mathcal{M}_2^{-1} \\ 0 & \mathcal{M}_2 + \mathcal{M}_1^{-1} & -\mathcal{M}_2 + \mathcal{M}_1^{-1} & 0 \\ 0 & \mathcal{M}_2 + \mathcal{M}_1^{-1} & \mathcal{M}_2 + \mathcal{M}_1^{-1} & 0 \\ \mathcal{M}_1 - \mathcal{M}_2^{-1} & 0 & 0 & \mathcal{M}_1 + \mathcal{M}_2^{-1} \end{pmatrix} \quad (39)$$

where $\mathcal{M}_1 = \frac{\omega_1 - \omega}{\omega_1 + \omega}$ and $\mathcal{M}_2 = \frac{\omega_2 - \omega}{\omega_2 + \omega}$ are the two diagonal elements of $\mathcal{M}(\omega)$ in (37). It can now be explicitly checked that $\widehat{\mathcal{M}}^{(B)}(\omega)$ and $\widehat{\mathcal{M}}^{(B=0)}(\omega)$ are related by the Ω matrix as was argued earlier. Furthermore, it is also straightforward to check that $\widehat{V}^{(B)}(x, t) = \Omega^T \widehat{V}^{(B=0)}(x, t) h(x)$ where

$$h(x) = \begin{pmatrix} \gamma & 0 & 0 & \delta \\ 0 & \gamma & -\delta & 0 \\ 0 & \delta & \gamma & 0 \\ -\delta & 0 & 0 & \gamma \end{pmatrix} \quad (40)$$

is independent of the spectral parameter as was pointed out earlier.

To summarize our results, it is important to note that we have revealed a deep connection between duality symmetry of two dimensional string effective action and its integrability properties through the construction of the monodromy matrix for the general case. Furthermore, we have shown that $\widehat{\mathcal{M}}(\omega)$ transforms nontrivially under the $O(d, d)$ transformations. Our results are likely to have applications in a variety of problems in string theory which are described by an effective two dimensional action (such as collisions of plane waves in the study of blackholes, cosmological scenarios etc). We have presented an application of our results in the study of an exactly solvable WZW model [14], which unravels some of the salient features of our investigation. As is evident, a wide class of problems can be studied from this perspective.

Acknowledgment: One of us (JM) acknowledges the warm hospitality of Prof. Y. Kitazawa and KEK. This work is supported in part by US DOE Grant No. DE-FG 02-91ER40685.

-
- [1] A. Giveon, M. Porrati and E. Rabinovici, Phys. Rep. **C244** (1994) 77.
 - [2] A. Sen, Developments in Superstring Theory, hep-th/9810044.
 - [3] J. Maharana, Recent Developments in String Theory, hep-th/9911200
 - [4] I. Bakas, Nucl. Phys. **B428** (1994) 374.
 - [5] J. Maharana, Phys. Rev. Lett. **75** (1995) 205; Mod. Phys. Lett. **A11** (1996).
 - [6] J. H. Schwarz, Nucl. Phys. **B447** (1995) 137, Classical Duality Symmetries in Two Dimensions, hep-th/9505170.
 - [7] A. Sen Nucl. Phys. **B447** (1995) 62 .
 - [8] P. Breitenlohner and D. Maison, Inst. H. Poincare, **46** (1987) 215; F. J. Ernst, A. Garcia and I. Hauser, J. Math. Phys. **28** (1987) 2155.
 - [9] H. Nicolai, Phys. Lett. **B235** (1990) 195.
 - [10] V. Belinskii and V. Zakharov, Sov. Phys. JETP, **48** (1978) 985; H. Nicolai, Schladming Lectures, Springer-Verlag, Berlin (1991) eds. H. Mitter and H. Gausterer.
 - [11] H. Nicolai, D. Korotkin and H. Samtleben, Integrable Classical and quantum Gravity, NATO Advanced Study Institute on quantum Fields and quantum spacetime, Cargese, 1996; hep-th/9612065.
 - [12] A. Das, "Integrable Models", World Scientific, Singapore (1989).
 - [13] A. Das, J. Maharana and A. Melikyan, Phys. Lett. **B518** (2001) 306, hep-th/0107229.
 - [14] C. Nappi and E. Witten, Phys. Lett. **B293** (1992) 309, hep-th/9206078.
 - [15] J. Maharana and J. H. Schwarz, Nucl. Phys. **B390**, hep-th/9207016.
 - [16] S. Mizoguchi, Nucl. Phys. **B461** (1996), A. Biswas, A. Kumar and K. Ray, Nucl. Phys. **B453** (1995) 181, A. A. Kehagias, Phys. Lett. **B360** (1995) 19, E. Abdalla and M. C. B. Abdalla, Phys. Lett. **B365** (1996) 41.
 - [17] M. Gasperini, J. Maharana and G. Veneziano, Phys. Lett. **B296** (1993) 51, hep-th/9209052.
 - [18] V. Bozza and G. Veneziano, JHEP **0010** (2000) 035, hep-th/0007159.